


Free Executive Summary



Review of the U.S. Army's Health Risk Assessments for Oral Exposure to Six Chemical-Warfare Agents

Subcommittee on Chronic Reference Doses for Selected Chemical Warfare Agents, National Research Council

ISBN: 0-309-06598-4, 320 pages, 6 x 9, paperback (1999)

This free executive summary is provided by the National Academies as part of our mission to educate the world on issues of science, engineering, and health. If you are interested in reading the full book, please visit us online at <http://www.nap.edu/catalog/9644.html>. You may browse and search the full, authoritative version for free; you may also purchase a print or electronic version of the book. If you have questions or just want more information about the books published by the National Academies Press, please contact our customer service department toll-free at 888-624-8373.

This executive summary plus thousands more available at www.nap.edu.

Copyright © National Academy of Sciences. Permission is granted for this material to be shared for noncommercial, educational purposes, provided that this notice appears on the reproduced materials, the Web address of the online, full authoritative version is retained, and copies are not altered. To disseminate otherwise or to republish requires written permission from the National Academies Press.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Summary

The U.S. ARMY is under a congressional mandate and the Chemical Weapons Convention of January 1993 to destroy its entire stockpile of chemical munitions. In addition to stockpiled munitions, nonstockpile chemical materiel (NSCM) has been identified for destruction. NSCM includes a host of lethal wastes from past disposal efforts, unserviceable munitions, chemically contaminated containers, chemical-production facilities, newly located chemical munitions, known sites containing substantial quantities of buried chemical weapons and wastes, and binary weapons and components. There are eight stockpile sites located in the continental United States and one on an island in the Pacific Ocean, and 82 NSCM locations have been identified. There are concerns, based on storage and past disposal practices, about soil and groundwater contamination at those sites. Six of the most commonly found chemical-warfare agents at stockpile and NSCM sites are the nerve agents GA, GB, GD, and VX and the vesicating (blistering) agents sulfur mustard and lewisite.

To ensure that chemical contamination is reduced to safe concentrations at stockpile and NSCM sites before they are used for residential, occupational, or wildlife purposes, the U.S. Army requested that health-based exposure limits for GA, GB, GD, VX, sulfur mustard, and lewisite be developed to protect the public and the environment. Oak Ridge National Laboratory (ORNL) was asked to conduct the health risk assessments and propose chronic oral reference doses (RfDs) and, where

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

appropriate, oral slope factors (SFs) for the six agents. RfDs are toxicological values developed for noncancer effects and used as reference points to limit human oral exposure to potentially hazardous concentrations of chemicals thought to have thresholds for their effects. RfDs are estimates (with uncertainty spanning an order of magnitude or greater) of daily oral chemical exposures that are unlikely to have deleterious effects during a human lifetime. For chemicals identified as carcinogens (e.g., sulfur mustard), SFs are also calculated. SFs are estimates of upper-bound lifetime cancer risk from chronic exposure to an agent.

The Army's Surgeon General adopted the proposed RfDs and SFs developed by ORNL as interim values to ensure that consistent health-based criteria were applied in ongoing initiatives requiring decisions on the safety of contaminated sites. The Army's Surgeon General also requested that the NRC independently review the scientific validity of these values. The NRC assigned this task to the Committee on Toxicology (COT), and a multidisciplinary subcommittee of experts was convened to assess the scientific validity of the interim RfDs developed for GA, GB, GD, VX, sulfur mustard, and lewisite and the SF developed for sulfur mustard. Specifically, the subcommittee was asked to (1) determine whether all the relevant toxicity data were considered appropriately; (2) review the uncertainty, variability, and quality of the data; (3) determine the appropriateness of the assumptions used to derive the RfDs (e.g., the application of uncertainty factors); and (4) identify data gaps and make recommendations for future research.

Although multiple agents are present at stockpile and NSCM sites, the subcommittee was asked to evaluate the agents only on an individual basis. Furthermore, although the most likely routes of exposure to chemical-warfare agents at these sites are the inhalation and dermal routes, the subcommittee was only asked to evaluate toxicological risk from the oral route at this time. The Army is in the process of developing inhalation exposure guidelines. The subcommittee was also not asked to address issues related to risk management, such as technology, detection, and feasibility.

EVALUATION OF THE ARMY'S INTERIM RFDS AND SFs

Table S-1 presents the interim RfDs and SFs adopted by the Army for GA, GB, GD, VX, sulfur mustard, and lewisite, as well as the recommenda

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

tions of the subcommittee. The subcommittee found that the guidelines used to derive the Army's interim RfDs were consistent with guidelines used by the U.S. Environmental Protection Agency (EPA) and were appropriate. In general, the approach was to identify the no-observed-adverse-effect level (NOAEL) or the lowest-observed-adverse-effect level (LOAEL) from animal or human studies. The NOAEL or LOAEL was divided by an overall uncertainty factor that reflects the uncertainties associated with the types of data used and a professional judgment of the entire data base for the chemical. An SF for sulfur mustard was derived using a comparative potency method.

TABLE S-1 Reference Doses and Slope Factors for Six Chemical-Warfare Agents

Agent	Army's Interim RfDs (mg/kg/d)	NRC's Recommended RfDs (mg/kg/d)	Army's Interim SFs (per mg/kg/d)	NRC's Recommended SFs (per mg/kg/d)
GA	4×10^{-5}	4×10^{-5}	NA	NA
GB	2×10^{-5}	2×10^{-5}	NA	NA
GD	4×10^{-6}	4×10^{-6}	NA	NA
VX	6×10^{-7}	5×10^{-7}	NA	NA
Sulfur mustard	7×10^{-6}	7×10^{-6}	9.5	1.6
Lewisite	1×10^{-4}	1×10^{-5}	NA	NA

Abbreviations: RfDs, reference doses; SFs, slope factors; NA, not applicable.

The subcommittee determined that the Army's interim RfDs for GA, GB, GD, and sulfur mustard were scientifically valid but concluded that the RfDs for VX and lewisite and the SF for sulfur mustard were too high. The bases for those conclusions are described below. Research recommendations for filling major data gaps are also presented.

CONCLUSIONS AND RECOMMENDATIONS

GA

The Army's interim RfD of 4×10^{-5} mg/kg of body weight per day for GA was based on a subchronic intraperitoneal toxicity study in rats, in which depression in plasma-cholinesterase (ChE) activity was considered the

critical end point. Although that end point is considered a biomarker of exposure rather than an adverse effect, the subcommittee agrees that the study is the best available one to use for deriving the RfD for GA and concludes that the available data on GA support the proposed RfD.

The major gap in the available information on GA is the lack of either a subchronic or a chronic oral toxicity study from which to derive the RfD. The absence of oral data could be addressed by conducting a subchronic oral toxicity study that assesses anti-ChE activity in red blood cells (RBCs) and plasma in one or preferably two species. If further research reveals that significant toxic effects can be induced by any of the nerve agents at doses below those that cause significant ChE inhibition, additional studies should be conducted to reassess the safety of the recommended RfD for GA.

GB

The Army's interim RfD of 2×10^{-5} mg/kg per day for GB was based on a subchronic oral toxicity study in rats, in which depression in RBC-ChE activity was considered the critical end point. Although that end point is a biomarker of exposure rather than an adverse effect, the subcommittee believes that this study is the best available one from which to derive the RfD for GB and concludes that the proposed RfD is scientifically valid.

The major gap in the available information on GB is the lack of either a subchronic or a chronic oral toxicity study that demonstrates a clear LOAEL or NOAEL. The absence of that type of data could be addressed by conducting a subchronic oral toxicity study that assesses anti-ChE activity in RBCs and plasma in one or preferably two species. If further research reveals that significant toxic effects can be induced by any of the nerve agents at doses below those that cause significant ChE inhibition, additional studies should be conducted to reassess the safety of the recommended RfD for GB.

GD

The Army's interim RfD of 4×10^{-6} mg/kg per day for GD was based on a subchronic oral toxicity study in rats, in which depression of plasma-

ChE activity was observed. Although that end point is a biomarker of exposure rather than an adverse effect, the subcommittee believes that this study is the best available one from which to derive the RfD for GD and concludes that the proposed RfD is scientifically valid.

The major gap in the available information on GD is the lack of either a subchronic or a chronic oral toxicity study that demonstrates a clear dose-response relationship between GD exposure and ChE inhibition. The absence of that type of data could be addressed by conducting a subchronic oral toxicity study that assesses anti-ChE activity in RBCs and plasma in one or preferably two species. Range-finding studies focusing on ChE analytical methods offer the best possibility for filling the data gap. If further research reveals that significant toxic effects can be induced by any of the nerve agents at doses below those that cause significant ChE inhibition, additional studies should be conducted to reassess the safety of the recommended RfD for GD.

VX

The Army's interim RfD of 6×10^{-7} mg/kg per day for VX was based on an oral toxicity study in sheep, in which depression in blood-ChE activity was observed. After evaluating that study, the subcommittee concludes that uncertainties about the relevance of this animal model to humans and weaknesses in the study design undermine the use of the study for deriving the RfD. Instead, the subcommittee recommends using a 1964 study of human volunteers in whom depression in RBC ChE was observed after oral exposure to low concentrations of VX. Although that study also has weaknesses and involves a biomarker of exposure rather than an adverse effect, the subcommittee believes it is preferable to use human data rather than data from a questionable animal model, because the uncertainty associated with extrapolating from animals to humans is avoided. On the basis of the human study, the subcommittee concludes that the data on VX support an RfD of 5×10^{-7} mg/kg per day, which is slightly lower than the Army's interim RfD of 6×10^{-7} mg/kg per day.

The major gap in the available information on VX is the lack of either a subchronic or a chronic oral toxicity study that demonstrates a clear dose-response relationship between VX exposure and ChE inhibition. The absence of that type of data could be addressed by conducting a

subchronic oral toxicity study that assesses anti-ChE activity in RBCs and plasma in one or preferably two species. If further research reveals that significant toxic effects can be induced by any of the nerve agents at doses below those that cause significant ChE inhibition, additional studies should be conducted to reassess the safety of the recommended RfD for VX.

SULFUR MUSTARD

The Army's interim RfD of 7×10^{-6} mg/kg per day for sulfur mustard was based on an oral two-generation reproductive toxicity study in rats, in which thickening of the forestomach epithelium was observed. The subcommittee agrees that this study is the best available one from which to derive the RfD for sulfur mustard and concludes that the interim RfD for sulfur mustard is scientifically valid. However, the subcommittee recommends adjustments in two of the uncertainty factors used to derive that RfD. Although the adjustments do not change the RfD for sulfur mustard, the subcommittee believes that they are scientifically justified and should be reflected in the Army's supporting documentation for the RfD.

Sulfur mustard is the only agent in this report associated with sufficient evidence of carcinogenicity in animal studies and, therefore, is the only agent for which a carcinogenic SF was derived. The indirect approach used to estimate the SF involved comparing the carcinogenic potency of sulfur mustard to that of the well-known carcinogen benzo[α]pyrene (B[α]P). Although the subcommittee finds that approach to be scientifically valid, given the absence of either a epidemiological investigation or a chronic oral animal bioassay on sulfur mustard, it recommends the use of a more recent risk estimate of the carcinogenic potency of B[α]P. On the basis of that estimate, the subcommittee concludes that the Army's interim SF of 9.5 per milligram per kilogram per day should be lowered to 1.6 per milligram per kilogram per day. Thus, if the potential carcinogenic risk from ingestion of sulfur mustard is restricted to less than 1 in 100,000 persons, daily oral doses should be limited to 6×10^{-6} mg/kg per day, a value slightly lower than the Army's interim RfD of 7×10^{-6} mg/kg per day.

The major gap in the available information on sulfur mustard is the lack of a chronic oral animal bioassay from which to derive the RfD and

SF. Because of that deficiency, the RfD for sulfur mustard is estimated by extrapolating from a subchronic study in animals, and the SF is established by applying comparative carcinogenic potency methods. The absence of chronic oral toxicity data can be addressed by conducting a chronic oral animal bioassay. It is important that sulfur mustard be delivered to animals at a slow rate (i.e., in the diet) rather than by stomach tube, because it is corrosive at the point of entry.

LEWISITE

The Army's interim RfD of 1×10^{-4} mg/kg per day for lewisite was based on two oral studies: a two-generation reproductive study and a 90-day toxicity study in rats. In both studies, necrosis and hyperplasia of the forestomach were observed. After considering those studies and other potential studies, the subcommittee concludes that a 1987 teratogenicity study conducted in rabbits is more appropriate than the rat studies for deriving the RfD, because there is evidence that the rabbit might be more susceptible to lewisite than the rat. On the basis of the rabbit study, in which maternal mortality and gastric lesions were observed, the subcommittee believes that the RfD for lewisite should be lowered from 1×10^{-4} mg/kg per day to 1×10^{-5} mg/kg per day.

The major gaps in the available information on lewisite are the lack of information on the implications of administering lewisite directly to the stomach over a short time and the absence of chronic oral toxicity data from which to derive an RfD. Because of those deficiencies, the RfD for lewisite was estimated by extrapolating from a less-than-ideal animal study to humans. Confidence in the RfD can be increased if subchronic oral toxicity studies in rabbits and rats are conducted to compare the effects of chronic oral exposure to low concentrations of lewisite with the effects of short-term intragastric administration of small volumes of lewisite. Such studies will provide not only the data needed to better understand the implications of dosing techniques but also more pertinent information on whether the rabbit is more appropriate than the rat for deriving an RfD for lewisite.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Review of the U.S. Army's Health Risk Assessments For Oral Exposure to Six Chemical-Warfare Agents

Subcommittee on Chronic Reference Doses For Selected Chemical-Warfare Agents
Committee on Toxicology
Board on Environmental Studies and Toxicology
Commission on Life Sciences
National Research Council

National Academy Press
Washington, D.C.

NATIONAL ACADEMY PRESS 2101 Constitution Ave., N.W. Washington, D.C. 20418

NOTICE: The project that is the subject of this report was approved by the Governing Board of the National Research Council, whose members are drawn from the councils of the National Academy of Sciences, the National Academy of Engineering, and the Institute of Medicine. The members of the committee responsible for the report were chosen for their special competences and with regard for appropriate balance.

The National Academy of Sciences is a private, nonprofit, self-perpetuating society of distinguished scholars engaged in scientific and engineering research, dedicated to the furtherance of science and technology and to their use for the general welfare. Upon the authority of the charter granted to it by the Congress in 1863, the Academy has a mandate that requires it to advise the federal government on scientific and technical matters. Dr. Bruce Alberts is president of the National Academy of Sciences.

The National Academy of Engineering was established in 1964, under the charter of the National Academy of Sciences, as a parallel organization of outstanding engineers. It is autonomous in its administration and in the selection of its members, sharing with the National Academy of Sciences the responsibility for advising the federal government. The National Academy of Engineering also sponsors engineering programs aimed at meeting national needs, encourages education and research, and recognizes the superior achievements of engineers. Dr. William A. Wulf is president of the National Academy of Engineering.

The Institute of Medicine was established in 1970 by the National Academy of Sciences to secure the services of eminent members of appropriate professions in the examination of policy matters pertaining to the health of the public. The Institute acts under the responsibility given to the National Academy of Sciences by its congressional charter to be an adviser to the federal government and, upon its own initiative, to identify issues of medical care, research, and education. Dr. Kenneth I. Shine is president of the Institute of Medicine.

The National Research Council was organized by the National Academy of Sciences in 1916 to associate the broad community of science and technology with the Academy's purposes of furthering knowledge and advising the federal government. Functioning in accordance with general policies determined by the Academy, the Council has become the principal operating agency of both the National Academy of Sciences and the National Academy of Engineering in providing services to the government, the public, and the scientific and engineering communities. The Council is administered jointly by both Academies and the Institute of Medicine. Dr. Bruce M. Alberts and Dr. William A. Wulf are chairman and vice chairman, respectively, of the National Research Council.

This project was supported by Contract Nos. DAMD 17-89-C-9086 and DAMD 17-99-C-9049 between the National Academy of Sciences and U.S. Department of Defense. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the organizations or agencies that provided support for this project.

International Standard Book Number 0-309-06598-4
Additional copies of this report are available from:
National Academy Press
2101 Constitution Ave., NW
Box 285
Washington, DC 20055
800-624-6242
202-334-3313 (in the Washington metropolitan area)
<http://www.nap.edu>

Copyright 1999 by the National Academy of Sciences. All rights reserved.

Printed in the United States of America

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

SUBCOMMITTEE ON CHRONIC REFERENCE DOSES FOR SELECTED CHEMICAL-WARFARE AGENTS

ROBERT SNYDER (*Chair*), Rutgers University College of Pharmacy and Environmental and Occupational Health Sciences Institute, Piscataway, New Jersey
MOHAMED B. ABOU-DONIA, Duke University, Durham, North Carolina
EDSON X. ALBUQUERQUE, University of Maryland School of Medicine, Baltimore, Maryland
JEFFREY I. DANIELS, Lawrence Livermore National Laboratory, Livermore, California
DONALD E. GARDNER, Inhalation Toxicology Associates, Raleigh, North Carolina
DAVID W. GAYLOR, U.S. Food and Drug Administration, Jefferson, Arkansas
ROGENE F. HENDERSON, Lovelace Respiratory Research Institute, Albuquerque, New Mexico
JOHN T. JAMES, National Aeronautics and Space Administration, Houston, Texas
SANFORD S. LEFFINGWELL, HLM Consultants, Auburn, Georgia
JOSEPH J. SAADY, Commonwealth of Virginia, Richmond, Virginia
PETER S. SPENCER, Oregon Health Sciences University, Portland, Oregon
BERNARD M. WAGNER, Consultant, Short Hills, New Jersey
BARRY W. WILSON, University of California, Davis California

Staff

KULBIR S. BAKSHI, Project Director
SUSAN N.J. PANG, Program Officer
RUTH E. CROSSGROVE, Editor
LINDA V. LEONARD, Senior Project Assistant
STEPHANIE PARKER, Assistant Information Specialist
Sponsor: U.S. DEPARTMENT OF DEFENSE

COMMITTEE ON TOXICOLOGY

BAILUS WALKER, JR. (*Chair*), Howard University Medical Center, Washington, D.C.
MELVIN E. ANDERSEN, Colorado State University, Denver, Colorado
GERMAINE M. BUCK, State University of New York at Buffalo
GARY P. CARLSON, Purdue University, West Lafayette, Indiana
JACK H. DEAN, Sanofi Pharmaceuticals, Inc., Malverne, Pennsylvania
ROBERT E. FORSTER II, University of Pennsylvania, Philadelphia, Pennsylvania
PAUL M.D. FOSTER, Chemical Industry Institute of Toxicology, Research Triangle Park, North Carolina
DAVID W. GAYLOR, U.S. Food and Drug Administration, Jefferson, Arkansas
JUDITH A. GRAHAM, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina
SIDNEY GREEN, Howard University, Washington, D.C.
WILLIAM E. HALPERIN, National Institute for Occupational Safety and Health, Cincinnati, Ohio
CHARLES H. HOBBS, Lovelace Respiratory Research Institute and Lovelace Biomedical and Environmental
Research Institute, Albuquerque, New Mexico
FLORENCE K. KINOSHITA, Hercules Incorporated, Wilmington, Delaware
MICHAEL J. KOSNETT, University of Colorado Health Sciences Center, Denver, Colorado
MORTON LIPPMANN, New York University School of Medicine, Tuxedo, New York
THOMAS E. MCKONE, University of California, Berkeley, California
ERNEST E. MCCONNELL, ToxPath, Inc., Raleigh, North Carolina
DAVID H. MOORE, Battelle Memorial Institute, Bel Air, Maryland
GÜNTER OBERDÖRSTER, University of Rochester, Rochester, New York
JOHN L. O'DONOGHUE, Eastman Kodak Company, Rochester, New York
GEORGE M. RUSCH, Allied Signal, Inc., Morristown, New Jersey
MARY E. VORE, University of Kentucky, Lexington, Kentucky
ANNETTA P. WATSON, Oak Ridge National Laboratory, Oak Ridge, Tennessee

Staff

KULBIR S. BAKSHI, Program Director
SUSAN N.J. PANG, Program Officer
ABIGAIL STACK, Program Officer
RUTH E. CROSSGROVE, Publications Manager
KATHRINE J. IVERSON, Manager, Toxicology Information Center
LUCY V. FUSCO, Project Assistant
LEAH PROBST, Project Assistant
EVELYN SIMEON, Project Assistant

BOARD ON ENVIRONMENTAL STUDIES AND TOXICOLOGY

GORDON ORIANS (*Chair*), University of Washington, Seattle, Washington
DONALD MATTISON (*Vice Chair*), March of Dimes, White Plains, New York
DAVID ALLEN, University of Texas, Austin, Texas
INGRID C. BURKE, Colorado State University, Fort Collins, Colorado
WILLIAM L. CHAMEIDES, Georgia Institute of Technology, Atlanta, Georgia
JOHN DOULL, The University of Kansas Medical Center, Kansas City, Kansas
CHRISTOPHER B. FIELD, Carnegie Institute of Washington, Stanford, California
JOHN GERHART, University of California, Berkeley, California
J. PAUL GILMAN, Celera Genomics, Rockville, Maryland
BRUCE D. HAMMOCK, University of California, Davis, California
MARK HARWELL, University of Miami, Miami, Florida
ROGENE HENDERSON, Lovelace Respiratory Research Institute, Albuquerque, New Mexico
CAROL HENRY, Chemical Manufacturers Association, Arlington, Virginia
BARBARA HULKA, University of North Carolina, Chapel Hill, North Carolina
JAMES F. KITCHELL, University of Wisconsin, Madison, Wisconsin
DANIEL KREWSKI, University of Ottawa, Ottawa, Ontario
JAMES A. MACMAHON, Utah State University, Logan, Utah
MARIO J. MOLINA, Massachusetts Institute of Technology, Cambridge, Massachusetts
CHARLES O'MELIA, Johns Hopkins University, Baltimore, Maryland
WILLEM F. PASSCHIER, Health Council of the Netherlands
KIRK SMITH, University of California, Berkeley, California
MARGARET STRAND, Oppenheimer Wolff Donnelly & Bayh, LLP, Washington, D.C.
TERRY F. YOSIE, Chemical Manufacturers Association, Arlington, Virginia

Senior Staff

JAMES J. REISA, Director
DAVID J. POLICANSKY, Associate Director and Senior Program Director for Applied Ecology
CAROL A. MACZKA, Senior Program Director for Toxicology and Risk Assessment
RAYMOND A. WASSEL, Senior Program Director for Environmental Sciences and Engineering
KULBIR BAKSHI, Program Director for the Committee on Toxicology
LEE R. PAULSON, Program Director for Resource Management

COMMISSION ON LIFE SCIENCES

MICHAEL T. CLEGG (*Chair*), University of California, Riverside, California
PAUL BERG (*Vice Chair*), Stanford University, Stanford, California
FREDERICK R. ANDERSON, Cadwalader, Wickersham & Taft, Washington, D.C.
JOHN C. BAILAR III, University of Chicago, Chicago, Illinois
JOANNA BURGER, Rutgers University, Piscataway, New Jersey
SHARON L. DUNWOODY, University of Wisconsin, Madison, Wisconsin
DAVID EISENBERG, University of California, Los Angeles, California
JOHN EMMERSON, Portland, Oregon
NEAL FIRST, University of Wisconsin, Madison, Wisconsin
DAVID J. GALAS, Keck Graduate Institute of Applied Science, Claremont, California
DAVID V. GOEDEL, Tularik, Inc., South San Francisco, California
ARTURO GOMEZ-POMPA, University of California, Riverside, California
COREY S. GOODMAN, University of California, Berkeley, California
HENRY HEIKKINEN, University of Northern Colorado, Greeley, Colorado
BARBARA S. HULKA, University of North Carolina, Chapel Hill, North Carolina
HANS J. KENDE, Michigan State University, East Lansing, Michigan
CYNTHIA KENYON, University of California, San Francisco, California
MARGARET G. KIDWELL, University of Arizona, Tucson, Arizona
BRUCE R. LEVIN, Emory University, Atlanta, Georgia
OLGA F. LINARES, Smithsonian Tropical Research Institute, Miami, Florida
DAVID LIVINGSTON, Dana-Farber Cancer Institute, Boston, Massachusetts
DONALD R. MATTISON, March of Dimes, White Plains, New York
ELLIOT M. MEYEROWITZ, California Institute of Technology, Pasadena, California
ROBERT T. PAINE, University of Washington, Seattle, Washington
RONALD R. SEDEROFF, North Carolina State University, Raleigh, North Carolina
ROBERT R. SOKAL, State University of New York, Stony Brook, New York
CHARLES F. STEVENS, The Salk Institute, La Jolla, California
SHIRLEY M. TILGHMAN, Princeton University, Princeton, New Jersey
JOHN L. VANDEBERG, Southwest Foundation for Biomedical Research, San Antonio Texas
RAYMOND L. WHITE, University of Utah, Salt Lake City, Utah

Staff

WARREN R. MUIR, Executive Director
JACQUELINE K. PRINCE, Financial Officer
BARBARA B. SMITH, Administrative Associate
KIT W. LEE, Administrative Assistant

OTHER REPORTS OF THE BOARD ON ENVIRONMENTAL STUDIES AND TOXICOLOGY

Research Priorities for Airborne Particulate Matter: II. Evaluating Research Progress and Updating the Portfolio (1999)

Ozone-Forming Potential of Reformulated Gasoline (1999)

Risk-Based Waste Classification in California (1999)

Arsenic in Drinking Water (1999)

Research Priorities for Airborne Particulate Matter: I. Immediate Priorities and a Long-Range Research Portfolio (1998)

Brucellosis in the Greater Yellowstone Area (1998)

The National Research Council's Committee on Toxicology: The First 50 Years (1997)

Toxicologic Assessment of the Army's Zinc Cadmium Sulfide Dispersion Tests (1997)

Carcinogens and Anticarcinogens in the Human Diet (1996)

Upstream: Salmon and Society in the Pacific Northwest (1996)

Science and the Endangered Species Act (1995)

Wetlands: Characteristics and Boundaries (1995)

Biologic Markers (5 reports, 1989–1995)

Review of EPA's Environmental Monitoring and Assessment Program (3 reports, 1994–1995)

Science and Judgment in Risk Assessment (1994)

Ranking Hazardous Waste Sites for Remedial Action (1994)

Pesticides in the Diets of Infants and Children (1993)

Issues in Risk Assessment (1993)

Setting Priorities for Land Conservation (1993)

Protecting Visibility in National Parks and Wilderness Areas (1993)

Dolphins and the Tuna Industry (1992)

Hazardous Materials on the Public Lands (1992)

Science and the National Parks (1992)

Animals as Sentinels of Environmental Health Hazards (1991)

Assessment of the U.S. Outer Continental Shelf Environmental Studies Program, Volumes I–IV (1991–1993)

Human Exposure Assessment for Airborne Pollutants (1991)

Monitoring Human Tissues for Toxic Substances (1991)

Rethinking the Ozone Problem in Urban and Regional Air Pollution (1991)

Decline of the Sea Turtles (1990)

Copies of these reports may be ordered from

the National Academy Press

(800) 624-6242

(202) 334-3313

www.nap.edu

Other Reports of the Committee on Toxicology

Assessment of Exposure-Response Functions for Rocket-Emission Toxicants (1998)
Review of A Screening Level Risk Assessment for the Naval Air Facility at Atsugi, Japan (Letter Report) (1998)
Review of Acute Human-Toxicity Estimates for Selected Chemical-Warfare Agents (1997)
The National Research Council's Committee on Toxicology: The First 50 Years (1997)
Toxicologic Assessment of the Army's Zinc Cadmium Sulfide Dispersion Tests (1997)
Toxicologic Assessment of the Army's Zinc Cadmium Sulfide Dispersion Tests: Answers to Commonly Asked Questions (1997)
Toxicity of Military Smokes and Obscurants, Volume 1 (1997) and Volume 2 (1999)
Toxicity of Alternatives to Chlorofluorocarbons: HFC-134a and HCFC-123 (1996)
Permissible Exposure Levels for Selected Military Fuel Vapors (1996)
Spacecraft Maximum Allowable Concentrations for Selected Airborne Contaminants, Volume 1 (1994), Volume 2 (1996), Volume 3 (1996)
Nitrate and Nitrite in Drinking Water (1995)
Guidelines for Chemical Warfare Agents in Military Field Drinking Water (1995)
Review of the U.S. Naval Medical Research Institute's Toxicology Program (1994)
Health Effects of Permethrin-Impregnated Army Battle-Dress Uniforms (1994)
Health Effects of Ingested Fluoride (1993)
Guidelines for Developing Community Emergency Exposure Levels for Hazardous Substances (1993)
Guidelines for Developing Spacecraft Maximum Allowable Concentrations for Space Station Contaminants (1992)

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Preface

Several military bases contaminated with chemical-warfare agents as a result of storage and past disposal practices are slated to be closed pursuant to the Base Realignment and Closure Act. Before those military bases can be transferred to civilian use, contaminated soil and water must be cleaned to levels that are considered safe. To help make decisions on restoration required at contaminated sites and on the potential uses of the former military installations (e.g., for housing, occupational, or wildlife purposes), the U.S. Army developed interim chronic oral reference doses and, where appropriate, oral slope factors for six chemical-warfare agents that are likely to be encountered at contaminated sites. Similar information for inhalation exposure is under development.

In this report, the Subcommittee on Chronic Reference Doses for Selected Chemical-Warfare Agents of the National Research Council's (NRC's) Committee on Toxicology reviews the scientific validity of the Army's interim values for the six chemical-warfare agents—GA, GB, GD, VX, sulfur mustard, and lewisite. The NRC report is intended to be useful to the Army in making site-specific cleanup decisions.

This report has been reviewed in draft form by individuals chosen for their technical expertise and diverse perspectives in accordance with procedures approved by the NRC's Report Review Committee for reviewing NRC and Institute of Medicine reports. The purpose of that independent review was to provide candid and critical comments to assist the NRC in making the published report as sound as possible and to ensure

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

that the report meets institutional standards for objectivity, evidence, and responsiveness to the study charge. The review comments and draft manuscript remain confidential to protect the integrity of the deliberative process. We wish to thank the following individuals, who are neither officials nor employees of the NRC, for their participation in the review of this report: Joseph Borzelleca, Virginia Commonwealth University; John Doull, University of Kansas Medical Center; Ronald W. Estabrook (report review committee monitor), University of Texas Southwestern Medical Center; Florence Kinoshita, Hercules Inc.; Loren Koller (report review coordinator), Oregon State University; John O'Donoghue, Eastman Kodak Company; and Joseph Rodricks, Life Sciences Trust.

The individuals listed above have provided many constructive comments and suggestions. It must be emphasized, however, that responsibility for the final content of this report rests entirely with the authoring committee and the NRC.

We gratefully acknowledge Veronique Hauschild, Joe King, and Steve Kistner (all of the U.S. Army Center for Health Promotion and Preventive Medicine) and Dennis Opresko, Robert Ross, Annetta Watson, and Robert Young (all of Oak Ridge National Laboratory) for providing background information and for making presentations to the subcommittee.

We are grateful for the assistance of the NRC staff for preparing the report. Staff members who contributed to this effort are James J. Reisa, director of the Board on Environmental Studies and Toxicology; Carol A. Maczka, senior program director for toxicology and risk assessment; Ruth E. Crossgrove, editor; and Linda Leonard, senior project assistant. We especially wish to recognize the major contributions of the project director, Kulbir S. Bakshi, and the program officer, Susan N.J. Pang, who directed the preparation of the subcommittee's report. Their knowledge of the scientific and technical literature and their tireless effort to obtain information and to organize the subcommittee meetings and the report aided in the successful completion of the project.

Finally, we would like to thank all the members of the subcommittee for their dedicated efforts throughout the development of this report.

ROBERT SNYDER, Ph.D.
CHAIR, SUBCOMMITTEE ON CHRONIC REFERENCE DOSES
FOR SELECTED CHEMICAL-WARFARE AGENTS
BAILUS WALKER, JR., Ph.D., M.P.H.
CHAIR, COMMITTEE ON TOXICOLOGY

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Contents

	Summary	1
1	Introduction	9
	References	16
2	Derivation of Reference Doses	17
	Reference-Dose Calculation	17
	Benchmark Dose	19
	Uncertainty Factors	20
	Conclusions	22
	References	22
3	Evaluation of the Army's Interim Reference Dose for GA	24
	Derivation of the Army's Interim RfD	24
	Appropriateness of the Critical Study	25
	Appropriateness of the Critical End Point	27
	Appropriateness of Uncertainty Factors	29
	Weight and Strength of Evidence	32
	Conclusions	32
	Data Gaps and Research Recommendations	33
	References	33

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

CONTENTS		xii
4	Evaluation of the Army's Interim Reference Dose for GB	36
	Derivation of the Army's Interim RfD	36
	Appropriateness of the Critical Study	37
	Appropriateness of the Critical End Point	38
	Appropriateness of Uncertainty Factors	40
	Weight and Strength of Evidence	43
	Conclusions	43
	Data Gaps and Research Recommendations	44
	References	44
5	Evaluation of the Army's Interim Reference Dose for GD	48
	Derivation of the Army's Interim RfD	48
	Appropriateness of the Critical Study	49
	Appropriateness of the Critical End Point	50
	Appropriateness of Uncertainty Factors	51
	Weight and Strength of Evidence	55
	Conclusions	55
	Data Gaps and Research Recommendations	55
	References	55
6	Evaluation of the Army's Interim Reference Dose for VX	59
	Derivation of the Army's Interim RfD	59
	Appropriateness of the Critical Study	60
	Appropriateness of the Critical End Point	62
	Appropriateness of Uncertainty Factors	63
	Weight and Strength of Evidence	65
	Conclusions	66
	Data Gaps and Research Recommendations	66
	References	67
7	Evaluation of the Army's Interim Reference Dose and Slope Factor for Sulfur Mustard	70
	Evaluation of the Army's Interim RfD	71
	Evaluation of the Army's Interim Cancer Slope Factor	76
	Conclusions	80
	Data Gaps and Research Recommendations	80
	References	81

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

CONTENTS		xiii
8	Evaluation of the Army's Interim Reference Dose for Lewisite	83
	Derivation of the Army's Interim RfD	83
	Appropriateness of the Critical Study	85
	Appropriateness of the Critical End Point	87
	Appropriateness of Uncertainty Factors	88
	Weight and Strength of Evidence	89
	Conclusions	90
	Data Gaps and Research Recommendations	90
	References	91
	Glossary	93
Appendix A:	Health Risk Assessment for the Nerve Agent GA	97
Appendix B:	Health Risk Assessment for the Nerve Agent GB	131
Appendix C:	Health Risk Assessment for the Nerve Agent GD (Soman)	167
Appendix D:	Health Risk Assessment for the Nerve Agent VX	197
Appendix E:	Health Risk Assessment for Sulfur Mustard (HD)	235
Appendix F:	Health Risk Assessment for Lewisite	275
Appendix G:	Inhibition of Cholinesterases and an Evaluation of the Methods Used to Measure Cholinesterase Activity	295

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

CONTENTS	xiv
----------	-----

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.

Review of the U.S. Army's Health Risk Assessments For Oral Exposure to Six Chemical-Warfare Agents

About this PDF file: This new digital representation of the original work has been recomposed from XML files created from the original paper book, not from the original typesetting files. Page breaks are true to the original; line lengths, word breaks, heading styles, and other typesetting-specific formatting, however, cannot be retained, and some typographic errors may have been accidentally inserted. Please use the print version of this publication as the authoritative version for attribution.